ticoke™Thoracoscopy Set

matic grasping forceps with -tipped and wide-opening jaws ow careful and precise manipuof the lung during diagnostic or peutic thoracoscopy.



coscopy:

May avoid open thoracotomy
Allows evaluation of parenchyma,
pleura, and mediastinum
May allow more accurate, minimally invasive staging
Dramatically reduces both hospital and postoperative recovery

s-on demonstrations and training with rick K. Toy, M.D.

Cabot Medical Booth

Monday, October 21

9:30-10:30 a.m. 2:30-3:30 p.m.

Tuesday, October 22

9:30-10:30 a.m. 2:30-3:30 p.m.

Wednesday, October 23

9:30-10:30 a.m. 2:30-3:30 p.m.

estral File (TM) - v	er 4.02	FAMILY GROUP	RECORD	14 APR 1992	Page
					ď.
BAND: Dr. Powell DAV	IES (AFN:BLF2-BW)				
N: [1780]	PLACE:	,		LOS ORDINANC	E DATA
.:	PLACE:	•		8:	
D:	PLACE:		•	B:	
.:	PLACE:			59.	
::	PLACE:			SS:	
1	PLHOL.			33.	
HER:					
HER:					
ER WIVES:					
E: Esther DAVIES	(AFN:8LF2-C3)		;		
N: [1]784]	PLACE:				
.:	PLACE:			₿:	
D:	PLACE:			ŧ :	
.:	PLACE:		i i	S∲:	
HER:					
HER:					
ER HUSBANDS:			1 1 1		
CHILDREN					
NAME: Franklin Jud	son DAVIES (AFN:23D0-D	K)	,		
- BORN: 2 Aug 1806	PLACE: Swegatchi	e, St. Lawrence, NY		B: 2 Oct 1	1967
CHR.:	PLACE:	,		E: 6 Feb 1	
		n Mactack III		SP:	
DIED: 16 Aug 1888	PLACE: Charlesto	n, wastach, ut	*	37.	
BUR.:	PLACE:	alt.			
	ND (AFN:23D0-FQ) and T	utners		SS: 6 Mar i	067 69
MAR.:	PLACE:			55: 6 nar 1	
NAME:					
- 60AN:	PLACE:			8:	
CHR.:	PLACE:			ŧ:	
DIED:	PLACE:			S₽:	
8UR.:	PLACE:				
SPOUSE:	, Ende.				
	PLACE:			SS :	
MAR.:	PLHCC;				
NAME:				<u> </u>	
- BORN:	PLACE:			₿;	
CHR.:	PLACE:			ŧ :	
DIED:	PLACE:			SP:	
BUA.:	PLACE:	,		1	
SPOUSE:			•		
maR.:	PLACE:			S\$:	
				,	
NAME:	PLACE:			8:	
- 80AN:				Į.	
CHR.:	PLACE:			SP:	
6.1.76	PLACE:			3,	
DIED:	PLACE:			1	•
BUR.:				1	
BUP.: Spouse:	****			ad	
BUR.:	PLACE:			, s\$:	
BUP.: Spouse:	:::::::::::::::::::::::::::::::::::::::	 ptized E=Endowed	SS-Sealed to Spouse	S\$: SP=Sealed to Parent	:::::::::::::::::::::::::::::::::::::::

The leaflet explains that during therapy the reaction is expected to be unsightly in the treated areas. In some cases this may last for several weeks following cessation of treatment. Response to Efudex occurs in the following sequence: erythema, usually followed by vesiculation, erosion, ulceration, necrosis and epithelization. The leaflet points out that it is time to stop Efudex if open sores develop. Complete healing of lesions may not be evident for one to two months following cessation of therapy with Efudex.

We hope you will find these materials helpful in your practice. If you would like a supply of these patient aids, please return the enclosed postage-paid reply card. Also, please see the complete listing of adverse effects, which is included in the enclosed complete product information. Should you have any questions, please contact our Professional Services Department at 201-812-2000.